

Communication



A multistep approach to preventing wrong site surgery in skin lesion excision

Nabil Mopuri, Quentin Frew, Charles Yuen Yung Loh, Peter Dziejewski
 St. Andrew's Centre for Plastic Surgery and Burns, Broomfield Hospital, Chelmsford, UK

Correspondence: Nabil Mopuri
 St. Andrews Centre for Plastic Surgery and Burns, Broomfield Hospital, 7 Whitfield Link, Chelmsford, CM2 9QF, UK
 Tel: +44-1245-266002, Fax: +44-1245-516295, E-mail: nabimopuri@yahoo.co.uk

Received: December 29, 2020 • Revised: May 25, 2021 • Accepted: May 27, 2021
 pISSN: 2234-6163 • eISSN: 2234-6171
<https://doi.org/10.5999/aps.2021.00353> • Arch Plast Surg 2021;48:570-571
 Copyright © 2021 The Korean Society of Plastic and Reconstructive Surgeons
 This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Wrong site surgery (WSS) is a never event and preventable error. It is well documented in various surgical specialties like urology, orthopedics and general surgery. There however is no reported data of outcomes of WSS in plastic surgery in the United Kingdom. We conducted a survey to assess the current incidence, factors involved and current protocols in place for plastic surgery units in both the United Kingdom and Ireland. We also devised an algorithm that is used at our unit currently that aims to reduce the incidence of WSS over a period from 2016 to 2018.

Sixty-six plastic surgery services in the United Kingdom and Ireland were contacted. The survey included incidence of WSS, anatomical location of lesion, level of experience of surgeon involved, related factors, any system in place to prevent WSS and any suggestions by the respondents on additional ways to avoid these events. The survey was sent to clinical leads of all plastic surgery units in the United Kingdom and Ireland by email.

We received replies from 19 of 64 clinical leads (30% response rate). In 42% of units (8/19) had experienced WSS. The number of such cases in the last 5 years varied with 26% (5/19) reporting one case, 10% (2/19) two cases, and 5% (1/19) three cases. The grade of surgeon involved included 7 of 19 consultants, 2 of 19 specialty doctors, 1 of 19 registrars and 0 of 19 senior house officers. Factors influencing WSS included wrong site marking by surgeon 20% (4/19) and failure to check the notes 20% (4/19) were the commonest causes. Other factors included checklist failures 10% (2/19), transfer of patient care to other surgeons 10% (2/19), inaccurate theatre lists 5% (1/19), failure to mark 5% (1/19), and failure to discuss with the patient 5% (1/19). Systems described that were already in place in-

cluded: use of the World Health Organization (WHO) checklist, preoperative marking by operating surgeon, or team briefing prior to list and operation commencing.

The incidence of WSS has been estimated at one out of 112,994 procedures; however, the number of unreported cases is estimated to be higher [1]. We introduced a strict guideline/protocol at our institute. A surgical grid form (Fig. 1), developed at our institute, identifies the anatomical site of surgery on the form printed with a line dia-

Mid Essex Hospital Services NHS Trust
St Andrew's Surgical Skin Lesion Site Marking

Patient Name: _____
 Record Number: _____
 NHS Number: _____
 Date of Birth: / / _____
 Sex: Male/Female _____
 Consultant: _____ Date: _____
 DIAGNOSIS: _____ OPERATION: _____

Affix Sticker if Available

Check List in Clinic

	Yes	No/NA
1. Skin lesion site marking form filled	<input type="checkbox"/>	<input type="checkbox"/>
2. Polaroid photograph with marking	<input type="checkbox"/>	<input type="checkbox"/>
3. Confirm with hand held mirror	<input type="checkbox"/>	<input type="checkbox"/>
4. Measured from lesion to any benign lesion/land mark /scar (For trunk and limbs) <small>Details</small>	<input type="checkbox"/>	<input type="checkbox"/>
5. Photograph request form signed by patient	<input type="checkbox"/>	<input type="checkbox"/>

Check list in Anaesthetic Room

	Date
1. Indelible skin Marking	<input type="checkbox"/>
2. Confirm with patient /Family member/Guardian	<input type="checkbox"/>
3. Confirm with colleague (Theatre staff)	<input type="checkbox"/>

Signature of the surgeon _____ Signature of patient _____

Fig. 1. Wrong site surgery prevention sample sheet used by St. Andrew's Centre for Plastic Surgery and Burns for skin lesion excision. LUA, left upper arm; RUA, right upper arm; B, back; BU, buttock; LLA, left lower arm; RLA, right lower arm; LH, left hand; RH, right hand; LUL, left upper leg; RUL, right upper leg; LLL, left lower leg; RLL, right lower leg; Ch, chest; Ab, abdomen; RP, right palm; LP, left palm; LT, left; RT, right; NA, not applicable.

gram in which the lesion is marked. A checklist on the other side of the surgical grid form confirms whether a photograph is taken after consent from the patient, whether confirmed with the patient by mirror, confirmed with distance from anatomical points to the site of lesion. On the day of surgery, the following steps are taken: (1) the clinic letter review, (2) reviewing the notes, (3) checking the line diagram in the notes, (4) clinical photograph of the lesion, (5) surgical grid form (anatomical marking form), (6) confirming site of surgery with the patient, (7) marking the lesion with indelible ink, or (8) confirming with colleagues in the anesthetic room before patient being taken into theatre. With the above steps, we have drastically reduced our WSS rate at our unit and hope others may follow suit.

Notes

Conflict of interest

No potential conflict of interest relevant to this article was reported.

Author contribution

Conceptualization: N Mopuri, Q Frew, CYY Loh, P Dziewulski. Data curation: N Mopuri, Q Frew, CYY Loh. Formal analysis: CYY Loh, P Dziewulski. Methodology: N Mopuri, CYY Loh. Project administration: N Mopuri, CYY Loh, P Dziewulski. Visualization: N Mopuri, CYY Loh, P Dziewulski. Writing - original draft: N Mopuri, CYY Loh. Writing - review & editing: N Mopuri, Q Frew, CYY Loh.

ORCID

Nabil Mopuri	https://orcid.org/0000-0002-2320-2700
Quentin Frew	https://orcid.org/0000-0002-2492-1635
Charles Yuen Yung Loh	https://orcid.org/0000-0003-2646-0358
Peter Dziewulski	https://orcid.org/0000-0003-2258-3115

Reference

1. Ambe PC, Sommer B, Zirngibl H. Wrong site surgery: incidence, risk factors and prevention. *Chirurg* 2015;86:1034-40.